


"CHI VA PIANO, VA SANO E VA LONTANO"

ARRIVO IN DEA...





Uomo, 53 aa

FdR: sovrappeso/tabagismo attivo


Dispnea da sforzi lievi Edemi declivi

COSA FACCIO ORA??



- A) Fuggo a gambe levate
- B) Mi faccio coraggio e vado a visitare il paziente

PRESENTAZIONE





EE: Non leucocitosi

Hb 15,1 g/dl

D dimero 2050 microg/l


Creatinina 1 mg/dl

NT-proBNP 4506 > pg/ml

PRESENTAZIONE







COSA AVRESTE FATTO AL MIO POSTO?



- A. Centralizzato subito il pz GUCH?
- B. Cardiovertito il pz in PS dopo ETE?
- C. Ricoverato e stabilizzato il pz?

IN REPARTO

Terapia diuretica infusionale

Ciclo di Levosimendan

Rate Control

CHA2DS2 VASC 1 HAS BLED 0 → NAO

NT-proBNP 4506 pg/ml --> 806

Recommendations for the management of AF in patients with congenital heart disease

Recommendations	Classa	Level ^b
 Oral anticoagulation should be considered in all adult patients with intracardiac repair, cyanosis, Fontan palliation, or systemic right ventricle and a history of AF, AFL, or intra-atrial re-entrant tachycardia. 1273 In patients with AF and other congenital heart diseases, anticoagulation should be considered in the presence of one or more non-sex stroke risk factor(s). 1273 	lla	с


OAC should be considered for stroke prevention in AF patients with a CHA₂DS₂-VASc score of 1 in men or 2 in women. Treatment should be individualized based on net clinical benefit and consideration of patient values and preferences.^{338,378,380}

lla

В



CORONAROGRAFIA





PROCEDURE DIAGNOSTICHE CATETERISMO - FASE: DURANTE PROCEDURA.

Sist.	Diast.	Media	Telediast.	Sat. 02 %
6	5	5		78
		3		75
28	16		6	73
22	15	17		74
		13		
115	12		14	98
120	80			98
	28 22 115	6 5 28 16 22 15 115 12	6 5 5 5 3 28 16 22 15 17 13 13 115 12	6 5 5 3 3 28 16 6 22 15 17 13 14

1/min

1/min

Wood

Wood

5,0

4,0

18,23

0,87

ardiac Output - Fick:

esistenze sistemiche:

esistenze polmonari:

ardiac Output - Termodiluzione:

Cardiac Index:

Cardiac Index:

Resistenze sistemiche:


Resistenze polmonari:


0,42

DIMISSIONE



TERAPIA DOMICILIARE
Canrenone 50 mg
Dabigatran 150 mg BID
Furosemide 25 mg 2 cp + 1 cp
Metoprololo 100 mg BID
Pantoprazolo 20 mg
Valsartan 40 mg 1 cp







ABLAZIONE TC

POST ABLAZIONE TC









UN MESE DOPO...









SEI MESI DOPO...







UN ANNO DOPO...



